



Ordered and non-ordered non-isometric convex quadrilaterals inscribed in a regular n -gon

Nesrine BENYAHIA TANI¹, Zahra YAHI², Sadek BOUROUBI³

¹ Algiers 3 University,

2 Ahmed Waked Street, Dely Brahim, Algiers, Algeria.

² Abderahmane Mira University, Bejaia, Algeria.

³ USTHB, Faculty of Mathematics

P.B. 32 El-Alia, 16111, Bab Ezzouar, Algiers, Algeria.

tani.nesrine@univ-alger3.dz or benyahiatani@yahoo.fr,
zahrayahi@yahoo.fr,
sbouroubi@usthb.dz or bouroubis@yahoo.fr

Abstract: Using several arguments, some authors showed that the number of non-isometric triangles inscribed in a regular n -gon equals $\{n^2/12\}$, where $\{x\}$ is the nearest integer to x . In this paper, we take back the same problem, but concerning the number of ordered and non-ordered non-isometric convex quadrilaterals, for which we give simple closed formulas, using Partition Theory. The paper is complemented by a study of two further kinds of quadrilaterals called proper and improper non-isometric convex quadrilaterals, which allows to give a connecting formula between the number of triangles and ordered quadrilaterals, which can be considered as a new combinatorial interpretation of certain identity in Partition Theory.

Keywords: Ordered parallel machines, multipurpose machines, complexity, heuristic, branch and bound.

1 Introduction

In the 1938's, Norman Anning from university of Michigan proposed the following problem [6]: "From the vertices of a regular n -gon three are chosen to be the vertices of a triangle. How many essentially different possible triangles are there?". For any given positive integer $n \geq 3$, let $\Delta(n)$ denotes the number of such triangles.

Using a geometric argument, the solution proposed by J.S. Frame, from Brown university, shown that $\Delta(n) = \{n^2/12\}$, where $\{x\}$ is the nearest integer to x . After that, other solutions were proposed by some authors, such as F. C. Auluck, from Dyal Singh college [2].

In 1978 Richard H. Reis, from the Southeastern Massachusetts university posed the following natural general problem: *From the vertices of a regular n -gon k are chosen to be the vertices of a k -gon. How many incongruent convex k -gons are there?*

Let us first precise that two k -gons are considered congruent if they are coincided at the rotation of one relatively other along the n -gon and (or) by reflection of one of the k -gons relatively some cord, that what we call non-isometric k -gons.

For any given positive integers $2 \leq k \leq n$, let $R(n, k)$ denotes the number of such k -gons. In 1979 Hansraj Gupta [5] gave the solution of Reis's problem, using the Möbius inversion formula.

Theorem 1

$$R(n, k) = \frac{1}{2} \binom{\lfloor \frac{n-h_k}{2} \rfloor}{\lfloor \frac{k}{2} \rfloor} + \frac{1}{2k} \sum_{d/\gcd(n,k)} \varphi(d) \binom{\frac{n}{d} - 1}{\frac{k}{d} - 1},$$

where $h_k \equiv k \pmod{2}$ and $\varphi(n)$ the Euler function.

One can find the first values of $R(n, k)$ in the Online Encyclopedia of Integer Sequences (OEIS) [7] as [A004526](#) for $k = 2$, [A001399](#) for $k = 3$, [A005232](#) for $k = 4$ and [A032279](#) for $k = 5$.

The immediate consequence of both Gupta's and Frame's Theorems is the following identity:

$$\left\{ \frac{n^2}{12} \right\} = \frac{1}{2} \left\lfloor \frac{n-1}{2} \right\rfloor + \frac{1}{6} \binom{n-1}{2} + \frac{\chi(3/n)}{3},$$

where $\chi(3/n) = 1$ if $n \equiv 0 \pmod{3}$, 0 otherwise.

In 2004 V.S. Shevelev gave a short proof of Theorem 1, using a bijection between the set of convex polygons with the tops in the n -gon splitting points and the set of all (0,1)-configurations with the elements in these points [8].

The aim of this paper is to enumerate the number of two kinds of non-isometric convex

quadrilaterals, inscribed in a regular n -gon, the ordered ones which have the sequence of their sides's sizes ordered, denoted by $R_O(n, 4)$ and those which are non-ordered denoted by $R_{\overline{O}}(n, 4)$, using the Partition Theory. As an example, let us consider the following figure showing three quadrilaterals inscribed in a regular 12-gon, the first is not convex, the second is ordered while the third is not. Observe that the second quadrilateral generates $1+1+3+3$ as partition of 8 in four parts, that is why it is called ordered.




Figure 1

2 Notations and preliminaries

We denote by G_n a regular n -gon and by \mathbb{N} the set of nonnegative integers. The partition of $n \in \mathbb{N}$ into k parts is a tuple $\pi = (\pi_1, \dots, \pi_k) \in \mathbb{N}^k, k \in \mathbb{N}$, such that

$$n = \pi_1 + \dots + \pi_k, \quad 1 \leq \pi_1 \leq \dots \leq \pi_k,$$

where the nonnegative integers π_i are called parts. We denote the number of partitions of n into k parts by $p(n, k)$, the number of partitions of n into parts less than or equal to k by $P(n, k)$ and by $q(n, k)$ we denote the number of partitions of n into k distinct parts. We sometimes write a partition of n into k parts $\pi = (\pi_1^{f_1}, \dots, \pi_s^{f_s})$, where $\sum_{i=1}^s f_i = k$, the value of f_i is termed as frequency of the part π_i . Let $m \in \mathbb{N}, m \leq k$, we denote $c_m(n, k)$ the number of partitions of n into k parts $\pi = (\pi_1^{f_1}, \dots, \pi_s^{f_s})$ for which $1 \leq f_i \leq m$ and $f_j = m$ for at least one $j \in \{1, \dots, s\}$. For example $c_2(12, 4) = 10$, the such partitions are 1128, 1137, 1146, 1155, 1227, 1335, 1344, 2235, 2244, 2334. Let $\delta(n) \equiv n \pmod{2}$, so $\delta(n) = 1$ or 0 , $\lfloor x \rfloor$ the integer part of x and finally $\{x\}$ the nearest integer to x .

3 Main results

In this section we give the explicit formulas of $R_O(n, 4)$ and $R_{\overline{O}}(n, 4)$.

Theorem 2 For $n \geq 4$,

$$R_O(n, 4) = \left\{ \frac{n^3}{144} + \frac{n^2}{48} - \frac{n\delta(n)}{16} \right\}.$$

Proof. First of all, notice that

$$R_O(n, 4) = p(n, 4). \quad (1)$$

Indeed, each ordered convex quadrilateral $ABCD$ inscribed in G_n can be viewed as a quadruplet of integers (x, y, z, t) , abbreviated for convenience, as a word $xyzt$, such that:

$$\begin{cases} n - 4 = x + y + z + t; \\ 0 \leq x \leq y \leq z \leq t, \end{cases} \quad (2)$$

where x , y , z and t represent the number of vertices between A and B , B and C , C and D and finally between D and A , respectively. It should be noted, that the number of solutions of System (2) equals $p(n, 4)$, by setting $x' = x + 1$, $y' = y + 1$, $z' = z + 1$ and $t' = t + 1$.

Now, let $g(z)$ be the known generating function of $p(n, 4)$ [3]:

$$g(z) = \frac{z^4}{(1-z)(1-z^2)(1-z^3)(1-z^4)}.$$

From expanding $g(z)$ in partial fractions, we obtain

$$g(z) = \frac{1}{32(1+z)^2} - \frac{13}{288(1-z)^2} - \frac{1}{24(1-z)^3} + \frac{1}{24(1-z)^4} + \frac{1-z^2}{8(1-z^4)} - \frac{1-z}{9(1-z^3)}.$$

Via straightforward calculations, it can be proved that

$$g(z) = \sum_{n \geq 0} \left(\frac{(-1)^n (n+1)}{32} - \frac{13(n+1)}{288} - \frac{(n+1)(n+2)}{48} + \frac{(1 + \frac{11}{6}n + n^2 + \frac{1}{6}n^3)}{24} + \epsilon(n) \right) z^n,$$

where $\epsilon(n) \in \{-\frac{17}{72}, -\frac{1}{8}, -\frac{1}{9}, -\frac{1}{72}, 0, \frac{1}{72}, \frac{1}{9}, \frac{1}{8}, \frac{17}{72}\}$.

Thus, we have

$$g(z) = \sum_{n \geq 0} \left(\frac{n^3}{144} + \frac{n^2}{48} + \frac{((-1)^n - 1)n}{32} + \beta(n) \right) z^n,$$

where $\beta(n) \in \{-\frac{5}{16}, -\frac{1}{4}, -\frac{29}{144}, -\frac{3}{16}, -\frac{5}{36}, -\frac{1}{8}, -\frac{13}{144}, -\frac{11}{144}, -\frac{1}{16}, -\frac{1}{36}, -\frac{1}{72}, 0, \frac{5}{144}, \frac{7}{144}, \frac{1}{9}, \frac{23}{144}, \frac{2}{9}, \frac{7}{72}\}$.

Since $p(n, 4)$ is an integer and $|\beta(n)| < 1/2$, we get

$$p(n, 4) = \left\{ \frac{n^3}{144} + \frac{n^2}{48} + \frac{((-1)^n - 1)n}{32} \right\}. \quad (3)$$

Hence, the result follows. ■

Remark 3 *G.E. Andrews and K. Eriksson said that the method used in the proof above dates back to Cayley and MacMahon [1, p. 58]. Using the same method [1, p. 60], they proved the following formula for $P(n, 4)$:*

$$P(n, 4) = \left\{ \frac{(n+1)(n^2 + 23n + 85)}{144} - \frac{(n+4) \lfloor \frac{n+1}{2} \rfloor}{8} \right\}.$$

Because $p(n, k) = P(n - k, k)$ (see for example [4]), it follows:

$$p(n, 4) = \left\{ \frac{n^3}{144} + \frac{n^2}{12} - \frac{n}{8} - \frac{n \lfloor \frac{n-1}{2} \rfloor}{8} \right\}. \quad (4)$$

Note that the formula (3) seems quite simple than (4).

To give an explicit formula for $R_{\overline{O}}(n, 4)$ we need the following lemma.

Lemma 4 For $n \geq 4$,

$$c_2(n, 4) = p(n, 4) - q(n, 4) - \left\lfloor \frac{n-1}{3} \right\rfloor.$$

Proof. By definition of $c_m(n, k)$ in section 2, it easily follows that

$$c_2(n, 4) = p(n, 4) - (q(n, 4) + c_3(n, 4) + \chi(4/n)),$$

where $\chi(4/n) = 1$ if $n \equiv 0 \pmod{4}$, 0 otherwise.

Furthermore, $c_3(n, 4)$ can be considered as the number of integer solutions of the equation:

$$3x + y = n, \text{ with } 1 \leq y \neq x \leq 1.$$

Since $x \neq y$, the solution $x = y = n/4$, when 4 divides n , must be removed. Then, by taking $y = 1$, one can get $c_3(n, 4) = \lfloor \frac{n-1}{3} \rfloor - \chi(4|n)$. This completes the proof. ■

Now we can derive the following theorem.

Theorem 5 For $n \geq 4$,

$$R_{\overline{O}}(n, 4) = \left\{ \frac{n^3}{144} + \frac{n^2}{48} - \frac{n\delta(n)}{16} \right\} + \left\{ \frac{(n-6)^3}{144} + \frac{(n-6)^2}{48} - \frac{(n-6)\delta(n)}{16} \right\} - \left\lfloor \frac{n-1}{3} \right\rfloor.$$

Proof. First of all, notice that $q(n, k) = p(n - k(k-1)/2, k)$ [1]. Then from (3) we get

$$q(n, 4) = p(n-6, 4) = \left\{ \frac{(n-6)^3}{144} + \frac{(n-6)^2}{48} - \frac{(n-6)\delta(n)}{16} \right\}.$$

Therefore, it is enough to prove that

$$R_{\overline{O}}(n, 4) = p(n, 4) + q(n, 4) - \left\lfloor \frac{n-1}{3} \right\rfloor. \quad (5)$$

In fact, each non-ordered convex quadrilateral may be obtained by permuting exactly two parts of some partitions of n into four parts, which is associated from System (2) to a

unique ordered convex quadrilateral. For example, in Figure 1 above, the ordered convex quadrilateral (b) assimilated to the solution 1133 of 8 or to the partition 2244 of 12, generates the non-ordered convex quadrilateral (c) via the permutation 1313. Obviously, not every partition of n can generate a non-ordered convex quadrilateral, those having three equal parts or four equal parts cannot. Also, each partition of n into four distinct parts $xyzt$ generates two non-ordered convex quadrilaterals, each one corresponds to one of the two following permutations $xytz$ and $xzyt$. On the other hand, each partition of n into two equal parts, like $xxyz$, with y and z both of them $\neq x$, generates only one non-ordered convex quadrilateral, corresponding to the unique permutation $xyxz$. Thus,

$$R_{\overline{O}}(n, 4) = 2q(n, 4) + c_2(n, 4), \quad (6)$$

Hence, from Lemma 4 the theorem holds. ■

Remark 6 *By substituting $k = 4$ in Theorem 1, we get*

$$R(n, 4) = \frac{1}{2} \binom{\lfloor \frac{n}{2} \rfloor}{2} + \frac{1}{8} \binom{n-1}{3} + \frac{n(1-\delta(n))}{16} + \alpha,$$

where

$$\alpha = \begin{cases} \frac{1}{8} & \text{if } n \equiv 0 \pmod{4}, \\ -\frac{1}{8} & \text{if } n \equiv 2 \pmod{4}, \\ 0 & \text{otherwise.} \end{cases}$$

Knowing furthermore that

$$R(n, 4) = R_O(n, 4) + R_{\overline{O}}(n, 4),$$

the following identity takes place according to Theorem 1 and Theorem 5:

$$\begin{aligned} \frac{1}{2} \binom{\lfloor \frac{n}{2} \rfloor}{2} + \frac{1}{8} \binom{n-1}{3} + \frac{n(1-\delta(n))}{16} + \alpha &= 2 \left\{ \frac{n^3}{144} + \frac{n^2}{48} - \frac{n\delta(n)}{16} \right\} + \\ &+ \left\{ \frac{(n-6)^3}{144} + \frac{(n-6)^2}{48} - \frac{(n-6)\delta(n)}{16} \right\} - \\ &- \left\lfloor \frac{n-1}{3} \right\rfloor. \end{aligned}$$

4 Connecting formula between $\Delta(n)$ and $R_O(n, 4)$

There are two further kinds of quadrilaterals inscribed in G_n , the proper ones, those which do not use the sides of G_n and the improper ones, those using them. In Figure 2 bellow,

two quadrilaterals inscribed in G_{12} are shown, the first one is proper while the second is not.




Figure 2

Let denote by $R_O^P(n, 4)$ and $R_O^{\bar{P}}(n, 4)$ respectively, the number of these two kinds of quadrilaterals. The goal of this section is to prove the following theorem.

Theorem 7 For $n \geq 4$,

$$\Delta(n) = R_O(n + 1, 4) - R_O(n - 3, 4).$$

Proof. Note first that an improper ordered quadrilateral is formed by at least one side of G_n , then the concatenation of the vertices of one of such sides gives a triangle inscribed in G_{n-1} , as shown in Figure 3.




Figure 3

Then we have

$$R_O^{\bar{P}}(n, 4) = \Delta(n - 1).$$

On the other hand, it is obvious to see that

$$R_O^P(n, 4) = p(n - 4, 4).$$

Then from (1), we get

$$R_O^P(n, 4) = R_O(n - 4, 4).$$

Since

$$R_O(n, 4) = R_O^P(n, 4) + R_O^{\bar{P}}(n, 4),$$

we obtain

$$R_O(n, 4) = R_O(n - 4, 4) + \Delta(n - 1).$$

So, the theorem has been proved while substituting n by $n + 1$. ■

Remark 8 *The well-known recurrence relation [4, p. 373],*

$$p(n, k) = p(n + 1, k + 1) - p(n - k, k + 1), \quad (7)$$

implies by setting $k = 3$,

$$p(n, 3) = p(n + 1, 4) - p(n - 3, 4). \quad (8)$$

Thus, as we can see, the formula of Theorem 7 can be considered as a combinatorial interpretation of identity (8).

For $k \leq n$, we have the following generalization, using the same arguments to prove Theorem 7.

Theorem 9 *For $n \geq k$,*

$$R_O(n, k) = R_O(n + 1, k + 1) - R_O(n - k, k + 1).$$

The formula of Theorem 9 can be considered as a combinatorial interpretation of the recurrence formula (7).

References

- [1] G. E. Andrews and K. Eriksson, '*Integer Partitions*', Cambridge University Press, Cambridge, United Kingdom, (2004).
- [2] N. Anning, J. S. Frame and F. C. Auluck, '*Problems for Solution: 3874-3899*', The American Mathematical Monthly, Vol. 47, No. 9, (Nov., 1940), 664-666.
- [3] A. Charalambos Charalambides, '*Enumerative Combinatorics*', Chapman & Hall /CRC, (2002).
- [4] L. Comtet, '*Advanced Combinatorics*', D. Reidel Publishing Company, Dordrecht-Holland, Boston, 1974, 133-175.

- [5] H. Gupta, Enumeration of incongruent cyclic k-gons, *Indian J. Pure Appl. Math.*, 10 (1979), no. 8, 964-999.
- [6] J. R. Musselman, G. Rutledge, N. Anning, W. Leighton, V. Thebault, '*Problems for Solution: 3891-3895*', *The American Mathematical Monthly*, Vol. 45, No. 9 (Nov., 1938), 631-632.
- [7] OEIS, The online Encyclopedia of Integer sequences, Published electronically at <http://www.research.att.com/~njas/sequences>, 2008.
- [8] V.S. Shevelev, Necklaces and convex k-gons, *Indian J. Pure Appl. Math.*, 35, no.5, (2004), 629-638.